

**FRUITS & VEGETABLES:** Mix 1½ fl. ozs. per 3 gals. of water (1 Tbsp./gal.) for a hand sprayer and apply to all plant foliage to cover an area of 1,000 square feet. Apply at first sign of infestation and repeat at weekly intervals when needed to maintain control of the following insects: **CABBAGE LOOPER** on Broccoli, Cauliflower, Collards, Kale, Mustard Greens, Turnip Greens, Cabbage, Celery, Lettuce, Melons and Tomatoes. **IMPORTED CABBAGE WORMS** on Broccoli, Cabbage, Cauliflower, Collards, Kale, Mustard Greens and Turnip Greens. **TOMATO HORNWORM** on Tomatoes.

### STORAGE AND DISPOSAL

**Do not contaminate water, food, or feed by storage or disposal.**

**STORAGE:** Store in original container in a cool, dry place inaccessible to children and pets and away from heat and direct sunlight. Protect from freezing. Storage at temperatures above 90°F may impair effectiveness.

**DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Do not reuse container. Wrap container and put in trash.

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed or inhaled. Avoid breathing vapors or spray mist. Avoid contact with skin, eyes or clothing. In case of contact immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

### STATEMENT OF PRACTICAL TREATMENT

**IF IN EYES:** Flush eyes with plenty of water. Get medical attention if irritation persists.

**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if irritation persists.

### ENVIRONMENTAL HAZARDS

Do not contaminate water when disposing of equipment washwaters.

Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith. Voluntary Purchasing Groups, Inc., warrants this product to conform to the chemical description on the label and for the purposes stated on the label. The limit of any liability shall be the purchase price paid by the user or buyer.



# Thuricide® Concentrate

- **Controls: Gypsy Moth, Cankerworm, Tent Caterpillar, Cabbage Looper, Imported Cabbage Worm & Tomato Hornworm**
- **Mix 4 Teaspoons per Gallon of Water for Shade Trees & Ornamentals.**

#### ACTIVE INGREDIENTS:

*Bacillus thuringiensis* subspecies *kurstaki*, Potency of 4,000 Spodoptera Units (at least 6 million viable spores per milligram\*)

Lepidopteran Active Toxin	0.8%
INERT INGREDIENTS**	99.2%
<b>TOTAL</b>	<b>100.0%</b>

\*Equivalent to 2.0 Billion International Units per Pint.

\*\*Contains Petroleum Hydrocarbon Solvent

Potency units should not be used to adjust use rates beyond those specified in the Directions for Use Section.

**KEEP OUT OF REACH OF CHILDREN CAUTION**  
See Side Panel for Additional Precautionary Statements

**NET CONTENTS ONE PINT (16 Fl.Ozs.)**

Distributed by  
Voluntary Purchasing Groups, Inc.  
Bonham, Texas 75418  
EPA Reg. No. 55947-134-7401  
EPA Est. No. 7401-TX-1  
Thuricide® is a Registered Trademark of Thermo Trilogy.  
31M-6-111K



### DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

Microbial Insecticide for the control of Certain Leaf-Chewing Worms on Shade Trees, Ornamentals, Fruits and Vegetables.

**SHAKE OR STIR BEFORE USING:** Pour recommended amount into nearly filled spray tank. Stir and add the balance of the water. Maintain agitation while spraying. Do not allow diluted spray to stand in tank for more than 12 hours. Good under leaf coverage is essential since the insect must digest THURICIDE®. THURICIDE® must remain on the leaf surface to be effective. Re-apply after heavy rains. Susceptible insects stop feeding after eating THURICIDE® but may otherwise appear to be unaffected for several days.

**SHADE TREES & ORNAMENTALS:** Mix 2 fl. ozs. per 3 gals. of water (4 tsp. per gal.) for a handsprayer or proportional amount for a hose-end sprayer following manufacturer's directions.

Apply at first sign of infestation and repeat at weekly intervals when needed to maintain control. Apply to thoroughly cover all foliage surfaces of shade trees and ornamentals for control of the following leaf-feeding worms:

Gypsy Moth (N.E.)*	Fall Cankerworm	Tent Caterpillar
Spring Cankerworm	Fall Webworm	

\*Apply when leaf expansion reaches 40% to 50%. If eggs hatch over a long period of time or if reinfestation occurs, spray about 14 days after first application.